

Fundamentals of Grape Integrated Disease Management for Beginners

Mizuho Nita
(Me-zoo-jo, or rhyme with Idaho or Navajo)
Associate Professor and Extension Grape Pathologist
Virginia Tech
AHS AREC at Winchester, VA
nita24@vt.edu

For Beginner's Workshop at AHS AREC
4 Nov 2021



COLLEGE OF AGRICULTURE AND LIFE SCIENCES
SCHOOL OF PLANT AND ENVIRONMENTAL SCIENCES
VIRGINIA TECH.



Outline

- ◆ Plant diseases and Integrated Pest Management
- ◆ Fungicide Resistance
- ◆ Pictures of common diseases
- ◆ Resources




Fungal diseases are very common in VA vineyards (or any vineyards located east of Rockies)

- ◆ Due to high humidity (rain and relative humidity) during the growing season
- ◆ Variety selection
 - ◆ Susceptible varieties such as 'Chardonnay' are preferred


A commercial vineyard in Loudoun county, VA



Grape diseases can be very serious!



It is not possible to grow susceptible variety without fungicides



How do these disease occur?

- Pathogens need to have a certain conditions to infect and cause disease



Host (grape)




**Environment
(esp. Rain & Temp)**



Black rot of grape



Pathogen




Fungicide resistance

- ◆ After several years of use, some of fungicides, especially newer ones, become less effective
- ◆ Many of new fungicides are targeting a specific gene or gene function
 - ◆ Highly specific and thus often safer to other organisms.
 - ◆ Use ounces not pounds
 - ◆ Other benefits such as movement of the chemical into plant tissues
 - ◆ In some cases, we can apply chemicals after a rain (typically, you need to apply before the rain)

The target pathogen can develop a resistance to the function = mode of action = how the pesticide kills or inactivates the target pathogen

- ◆ Pathogens can become less sensitive to a fungicide because...
 - ◆ Some of population (isolates) were not sensitive to begin with
 - ◆ Mutation of the target gene (or gene function) happened after exposed to the fungicide.



An example of fungicide resistance development QoI or Strobilurin fungicides

- ◆ The first fungicides in this family were isolated from wood-rotting mushroom fungi, including one called *Strobilurus tenacellus*.
- ◆ All QoI fungicides share a common biochemical mode of action:
 - ◆ Interfere with energy production in the fungal cell.
 - ◆ They block electron transfer at the site of quinol oxidation (the Qo site) in the cytochrome *bc₁* complex, thus preventing ATP formation.
- ◆ It has curative activity against some of pathogens = you can apply after infection takes place.

QoI fungicide was introduced in late 1990's, and it was working against multiple pathogens

- ◆ However, this entire group was found to be no longer effective against both grape downy and powdery mildew in VA by 2007-09
- ◆ Only 10-12 applications were enough for fungal pathogens to develop resistance to the QoI
 - ◆ A single mutation site was often associated with the resistant isolates
- ◆ Once developed, the resistance highly likely stay for good = you cannot use the same mode of action any longer.

Once developed, fungicide resistance will stay...



Best way to avoid fungicide resistance are tank mix, limitation of the use, and rotation of mode of action

- ◆ Some of fungicides are less prone to the development of resistance because they have multiple modes of action
 - ◆ **Sulfur** for powdery mildew, **mancozeb** for downy mildew, black rot, and Phomopsis, **copper** for downy and powdery mildew, and **captan** for downy mildew, Phomopsis and Botrytis bunch rot
 - ◆ Mixing them with a newer fungicide has shown some evidence of delaying onset of resistance with some of pathogens


You cannot use the same materials repeatedly

- ◆ Often time there is a legal limitation in number of applications or amount of the chemical you can use per season
 - ◆ Example with grape: Mancozeb's PHI (Pre-Harvest Interval) is 66-day, *plus* there is a limitation on the amount (19.2 lb of a.i./ acre/ season)
- ◆ Recommendations on new fungicides are to apply no more than two applications per season (listed on the label = legal)
- ◆ → Rotate with different modes of action!
 - ◆ **However, rotating product or chemical name may not result in the rotation of mode of action!**

I hope things are more straightforward, but it is not...


- ◆ **Different products may have the same mode of action**
 - ◆ Both 'Elite' and 'Orius' have a tebuconazole as an active ingredient (a.i.), and tebuconazole belong to a mode of action DMI (demethylation inhibitors, or also called sterol inhibitor or SI)
- ◆ **Different chemicals may have the same mode of action**
 - ◆ Both 'Elite' (a.i. = tebuconazole) and 'Rally' (a.i. = myclobutanil) belongs to DMI group

FRAC (Fungicide Resistant Action Committee) code
<http://www.frac.info/>



Sometimes, one product has two modes of action

- Increase efficacy
- May delay the onset of resistance
- Make it more difficult to rotate!



Rotation of fungicide alone is complicated enough!
 Please plan ahead!!

Proper planning will help you to:

- ◆ Prepare time and resources
 - ◆ Be thorough and realistic
- ◆ Check inventory of your supplies
- ◆ Remember what you did last year
 - ◆ Lower the risk of making the same mistake
- ◆ Recognize which diseases were more prevalent
 - ◆ = Adjustment for a challenging season!!



Let's go through common diseases that you probably will see in your vineyards!

- ◆ For the sake of time, I will focus major fungal diseases; however, there are diseases caused by
 - ◆ Viruses
 - ◆ leafroll viruses, red blotch, etc. (60+)
 - ◆ Bacteria
 - ◆ Pierce's Disease, crown gall
 - ◆ Phytoplasma
 - ◆ grapevine yellows



The correct identification is critical because different management tools will be needed for seemingly similar diseases



Downy Mildew

The infection conditions and chemical to be used are different!

Powdery Mildew



Downy Mildew

- ◆ Caused by Oomycete pathogen, *Plasmopara viticola*, which can infect leaves and berries, berry infection can cause serious damage
- ◆ Heavy leaf infection can cause a defoliation

Oily spot appearance on upper surface



Pictures taken from Organic grape production guide: OSU, Ellis and Nita 2004

Downy Mildew



Powdery Mildew

- ◆ Caused by a fungal pathogen, *Erysiphe necator* (= *Uncinula necator*)
- ◆ It can infect leaves and berries, berry infection can cause serious damage
- ◆ Infection of berries during early season can increase the risk of other diseases



Pictures taken from Organic grape production guide: OSU, Ellis and Nita 2004

Powdery Mildew



It can be found on the both upper and lower surface, but more commonly found on the upper surface

Phomopsis Cane and Leaf Spot

- ◆ Caused by a fungus, *Phomopsis viticola*.
- ◆ It can infect leaves, canes, rachis, and berries (up to 30% loss of yield has been reported), it can cause premature drop of berries
- ◆ Even though it does not cause major damage, it can cause a slow decline of vines




Phomopsis cane and leaf spot



Black Rot

- ◆ It is caused by a fungus, *Guignardia bidwellii* that can infect leaves and berries, berry infection can cause serious damage
- ◆ Infected berries will produce spores next year



Black Rot



Botrytis bunch rot, or gray mold

- ◆ It is caused by a fungus *Botrytis cinerea*.
- ◆ It can cause damage to berries, and can be very significant
- ◆ The gray moldy appearance is due to mass of conidia
- ◆ It has wide range of hosts, strawberry and other small fruits, crop debris, etc...





Botrytis Bunch Rot



How do we manage these diseases?

- ◆ Integrated Pest Management (IPM)
- ◆ Disease triangle matters!
 - ◆ You will see more warm season disease such as ripe rot and Pierce's Disease in southeastern part of the state
 - ◆ Grape changes its susceptibility to disease over the course of the season too!





Extension Resources

In-person Extension meetings
Please attend our grape disease and IPM workshops

Extension publications

Grapes: Diseases and Insects in Vineyards

Douglas G. Pfeiffer, Extension Entomologist, Virginia Tech
Anton B. Baudoin, Plant Pathologist, Virginia Tech
J. Christopher Bergh, Extension Entomologist, Alton H. Smith Jr. AREC
Mizuko Nita, Extension Plant Pathologist, Alton H. Smith Jr. AREC

Additional information on pest and beneficial species identification is available online at <http://www.virginiafruit.cento.vt.edu/>. Disease updates and management information is available at <http://www.grapepathology.blogspot.com>.

In January 2018, a new invasive insect was found in Virginia from expanding its range. More than 70 species of insects are known to cause injury on some. Application rates: T

Blog (grapepathology.blogspot.com)
Twitter @grapepathology
Facebook GrapePathVATech

Web-based applications

Resources on Grape Disease Management

- ◆ My blog
 - ◆ Updated frequently during the season
- ◆ **Ext.grapepathology.org**
- ◆ I will upload today's presentation!

VCE's Pest Management Guide (PMG)

- It covers not only diseases, but also insect and weeds
- Updated every year


Diseases and Insects in Vineyards
Douglas G. Pfeiffer, Extension Entomologist, Virginia Tech
Anton B. Baudoin, Plant Pathologist, Virginia Tech
J. Christopher Bergh, Extension Entomologist, Alison H. Smith Jr. AREC
Mizuno Nita, Extension Plant Pathologist, Alison H. Smith Jr. AREC

Additional information on pest and beneficial species identification is available online at <http://www.virginiafruit.ento.vt.edu/>.
 Application rates: The rate per acre column gives rates for low-volume or concentrate applications. Sprays may be applied as semi-concentrate (40-100 gal/A) or concentrate (10-40 gal/A) sprays. Use caution with more concentrated sprays; the smaller droplet sizes associated with low-volume application are more prone to drift. Amount of pesticide to be applied for dilute applications (usually 100 gal/A early in early season, 200 gal/A in mid season, and 300 gal/acre in late season) is usually given on the label.

Pest	Chemical and Formulation	Rate/Acre	Spray Timing and Remarks
Dormant			
Anthraxnose (Bird's eye rot), Powdery Mildew, Phomopsis	lime sulfur solution	10.0 gal	Only necessary where anthracnose, Phomopsis, or powdery mildew have been a serious problem. Lime sulfur can reduce overwintering inoculum of these diseases.
Mealybugs			
	Applaud 70DF	9.0-12.0 oz	If a problem at harvest in the previous year. If a delayed dormant spray does not provide a adequate control, a summer application may be made. Baythroid targets only crawlers. Movento prebloom only in table grapes.
	Venom 20SG	0.44-0.66 lb (foliar)	
		1.13-1.32 lb (soil)	
	Assail 30SG	2.5 oz	
	Provado Solupak	0.8-1.0 oz	
	Baythroid 2EC	2.4-3.2 fl oz	
	Movento 2SC	6.0-8.0 fl oz	
Bud Swell			
Grape flea beetle	Danitol 2.4EC or	8.0 fl oz	If adult beetles are present in damaging numbers. See Table 3.4 for Restricted Entry.

Grape disease management guide for non-bearing grapes

- Aiming for first- and second-year vines which should not bear fruits.



Virginia Cooperative Extension
Virginia Tech • Virginia State University
www.ext.vt.edu


Fungicide Spray Guidelines for Non-bearing Vineyards

Authored by Mizuho Nita, Associate Professor and Extension Grape Pathology Specialist, Alton H. Smith Jr. Agricultural Research and Extension Center, School of Plant and Environmental Sciences, Virginia Tech

Introduction
The approach to non-bearing vines is different from bearing vines because you do not need to protect berries. Also, some diseases such as Phomopsis cane and leaf blight tend to appear in the vineyards that are established for several years. This guide intends to provide examples to aid in building your spray program.



For vineyards in Virginia, we have to deal with multiple diseases due to our environmental conditions. The development of diseases depends on vineyard history, cultivar, proximity to the existing vineyards and wild grapes, weather conditions, and other factors. Thus, you need to adjust your spray program to account for all of these conditions. For example, I need to protect vines from powdery mildew for my Chardonnay vineyard from nearly the beginning to the end. Still, with my Cabernet Sauvignon vineyard, located less than 100 feet away, I can skip some powdery mildew sprays.

Seasonal Fungicide Spray Guideline for Non-bearing Vineyards

Growth stage or timing	Material and rate/Acre	Comments
New shoots The first spray Target diseases Phomopsis cane and leaf spot and downy mildew Timing Begin at ~ 3- to 5-inch shoot  Grape illustrations are adapted from Eichhorn and Lorenz, 1977	Option A mancozeb at 3 lb/A	In non-bearing vineyards (1st and 2nd year), you may use a simplified program to control black rot, Phomopsis, downy mildew, and powdery mildew. The main focus will be on downy and powdery mildew during the first year. A protection program starts when shoots are about 3 to 5 inches in length. The target disease is Phomopsis cane and leaf spot, which should not appear in a new vineyard, but it may happen if you have a vineyard nearby. Both mancozeb and captan control downy mildew. Powdery mildew is less likely active at this time of the season.
	Option B captan at 2 to 3 lb/A	
	Option C mancozeb at 3 lb/A	
12" – 18" shoots 2 nd and 3 rd sprays	Option A Fixed copper	For option A, fixed copper is listed as the first option for this spray, and mancozeb is listed as the second option. We need to compare

My Fungicide application workbooks

- With pictorial keys for the target host stage
- You should have a non-bearing vine version in the package
- We are working on an online version
- GrapeIPM.org

Growth stage or timing	Material and rate/acre	Comments
3- to 5-inch shoot or 7-10 days after the last spray 	Same as 1/2- to 1-inch shoot spray ----- Note ----- When you wonder about which fungicide to use, think about what was the primary problem in your vineyard, and also what is going on in this season. For example, if you had BR problems last year, and a weather forecast shows a warm rain event, you want to incorporate a fungicide against BR (such as Elite or Rally) and apply it before the rain.	Some of you start your program at this stage. Just remember that from 1-inch to 5-inch takes only a few days! Most fungicides act only as protectants. Thus, in order to protect new growth from fungal infection, these materials need to be applied before the rain. 7-day interval application needs to be considered if: <ul style="list-style-type: none"> • you are applying sulfur for PM (which does not require rain to infect tissue), • PM has been a concern in your vineyard • there has been a lot of rain since the last spray, or • it is unusually warm, and shoots are growing rapidly. If rain is predicted between 7 and 10 days after your last spray, make another application before the rain.
6- to 10-inch shoot or 7-10 days after the last spray 	Same as 1/2- to 1-inch shoot spray	Please see above. To lower risk of fungicide resistance development, rotate the mode of action. In general, 2-3 sprays of a resistance-prone fungicide (3 for SI and 2 for strobilurin) per season are the maximum recommended. Please plan ahead. Refer to Table 2 for the mode of action, and read and follow the label.

Southeastern Regional IPM guide

<http://www.smallfruits.org/SmallFruitsRegGuide/>

- ◆ Another version of the management guide
- ◆ Why so many? - there are many ways to look at pest management

2021 Southeast Regional Bunch Grape Integrated Management Guide

Commodity Editor
Mizuho Nita (Virginia Tech)

Section Editors

Pathology: Phil Brannen (University of Georgia), Sarah Villani (North Carolina State University), Zack Hansen (University of Tennessee), Rebecca Melanson (Mississippi State) and Mizuho Nita (Virginia Tech)
Entomology: Hannah Burrack (North Carolina State University), Doug Pfeiffer (Virginia Tech), Brett Blaauw (University of Georgia), and Frank Hale (University of Tennessee)
Weed Science: Wayne Mitchem (North Carolina State University)
Vertebrate Management and Grape Culture: David Lockwood (University of Tennessee); Michael T. Mengak (University of Georgia), and Eric T. Stafne (Mississippi State University)
Pesticide Stewardship and Safety: Ash Sial (University of Georgia)

Senior Editors
Phil Brannen (University of Georgia)
Bill Cline (North Carolina State University)

A product of the Southern Region Small Fruit Consortium (www.smallfruits.org). Recommendations are based on information from the manufacturer's label and performance data from research and extension field tests. Because environmental conditions and grower application methods vary widely, suggested use does not imply that performance of the pesticide will always conform to the safety and pest control standards indicated by experimental data. This publication is intended for use only as a guide. Specific rates and applications methods are on the pesticide label, and these are subject to change at any time. Always refer to and read the pesticide label before making any application! The pesticide label supersedes any information contained in this guide, and it is the legal document referenced for application standards.

Online resource demonstration

- ◆ <http://ext.grapepathology.org>
- ◆ <http://grapeIPM.org>
- ◆ <https://newa.cornell.edu/>

Acknowledgement

2019-21 Grape Path Team

Abdullah Nahiyen

Amanda Bly and Diana Scorpio

Morgan Gannon

Akiko Nita

Dr. Diana McHenry

Akiko Mangan

Robert Burgholzer

Kenny Savia

Redbubble.com